

Central and Northern California Coastal Marine Habitats: Oil Residence and Biological Sensitivity Indices

RELATIVE ABUNDANCE OF MAJOR MACROBIOTA*

Taxa	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S
ROCKY INTERTIDAL MACROBIOTA																			
Acorn barnacles		0		0	0	0	0	0		0							0		0
Chlorophyta		C		0	0	0	0	0		0									
Porphyra spp.		0		0	0	0	0	0					0		C	C	C		C
Pelvetiopsis limitata				0	0	0	0	0		0									0
Endocladia muricata		C		C	C	C	C	C		C					0	0	0		0
Pelvetia fastigiata				0	0	0	0	0		0					0	0	0		0
Fucus distichus															0	0	0		0
Gigartina spp.		0		C	C	C	C	C		C							C	C	A
Pollicipes polymerus		C		0-C	0-C	0-C	0-C	0-C		C					A	C	C	C	C
Mytilus californianus		C		C	C	C	C	C-A		C					A	C	C	C	C-A
Corallina spp./Gigartina spp.		C		C	0	0	0	0		C					C	C	C	C	C
Postelsia palmaeformis				C	C	C	C	C		C					C	C	C	C	C
Halosaccion glandiforme		0													0	0	0		C
Iridaea spp.				C	0	0	0	C		C					C	C	C	C	C
Odonthalia spp./Rhodomela larix																			C
Coralline algae				C	0	0	0	C		C					A	C	C	C	A
Phyllospadix spp.		0		C	C	0	0	C		C					C	C	C	C	0
Alaria marginata						C	C	C		C					C	C	C	C	0
Egregia menziesii		C		C				C		C					C	C	C	C	0
Laminaria spp.		0		C	0	0		C		C					C	C	C	C	0
Lessoniopsis littoralis				0											C	C	C	C	0
OTHER MACROBIOTA																			
Kelp beds:																			
Macrocystis spp.										0		0	0	0	0	0	0	0	C
Nereocystis luetkeana						0-C													
Marine mammals:																			
Elephant seal																			
Harbor seal						0		0				0-A	0-A		0		0-C		0-C
Steller sea lion															0				
California sea lion															0				0-C
Seabird nesting colonies:																			
Fork-tailed storm petrel																			
Leach's storm petrel																			
Ashy storm petrel																			
Brandt's cormorant								0											
Double-crested cormorant																			
Pelagic cormorant						0		A											
Black oystercatcher						0		0											
Western gull						0		0											
Common murre								A											
Pigeon guillemot								0-A											
Cassin's auklet						C		0-A											
Rhinoceros auklet																			
Tufted puffin																			
Threatened/Endangered Species:																			
Aleutian Canada goose																			
Southern sea otter																			

*Relative abundance for summer conditions: 0 = occasional, C = common, A = abundant

PHYSICAL SHORE-ZONE CHARACTERISTICS

UNIT IDENTIFIERS	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S
ALONGSHORE LENGTH (km)	0.4	0.9	1.7	0.6	1.4	0.6	1.6	0.8	0.5	0.3	0.4	1.3	1.9	0.2	0.8	0.5	1.6	0.5	0.0
ACROSS-SHORE WIDTH (m)	50	5-20	30-50	5-20	5-20	<5	5-20	<5	25	<5	25	30	5-20	<5	20-75	<10	15-30	20	-
WAVE EXPOSURE	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	1-2
ACROSS-SHORE COMPONENTS (morphology, texture)	Ca, Csp Bf, Csp	Ca, Rm Bi, Crb: Rm	Dl, Cs Bb, Csp Bf, Cs	Ca, Rm Bi, Crb: Rm	Ca, Rm Bi, Crb: Rm	Cc, Rm Ore, Rm	Ca, Rm Bi, Crb: Rm	Cc, Rm Ore, Rm	Ca, Rs Bi, Csp	Cc, Rm Ore, Rm	Ca, Rs Bi, Csp	Cp, Cg Bb, Cs Bf, Cs	Cc, Rs Pi, Rs Ore, Rs	Ca, Rs Pi, Rs Ore, Rs	Ca, Rs Bi, Cs Pf, Rs Ore, Rs	Ca, Rs Bi, Csp: Rs	Ca, Rs Bi, Csp: Rs	Cp, Rs Bf, Csp Oe, Rs	Eb, Cs Ar Is, Ar
Primary		Ca, Rm Bi, Cr Bi, Csg	As, Ar Bb, Csp Bf, Cs	Ca, Rm Bi, Cr Bi, Csg	Ca, Rm Bi, Cr Bi, Csg		Ca, Rm Bi, Cr Bi, Csg						Cc, Rs Pi, Ccb Rs Ore, Rs			Oe, Rs	Ca, Rs Bi, Cg Pf, Crb Rs	Cp, Rs Bi, Csp Pl, Rs	
Secondary																			
MICRO RELIEF	S	R	S	R	R	R	R	R	S	R	S	S	R	R	R	S	R	S	S
MACRO RELIEF	S	S	S	S	S	R	S	R	S	R	S	S	R	R	S	S	R	S	S-R
SUMMARY CHARACTERISTICS	bg	Rbmv	bg	Rbmv	Rbmv	R	Rbmv	R	bs	R	bs	bs	Rbmv	R	RPbs	bm	RPbm	RPbs	E
OIL RESIDENCE INDEX	3	3	2	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	1
GROUND TRUTH	0	3	3	0	0	0	0	0	0	0	0	3	2	0	0	2	3	0	0

* Significant seasonal beach accretion during summer.

ABBREVIATED PHYSICAL SHORE-ZONE CODING EXPLANATION

This is an abbreviated legend for the physical shore-zone coding sheets; consult the main text for a complete discussion of codes and rationale. Only the across-shore descriptors for morphology and texture are discussed.

ACROSS-SHORE COMPONENTS - dominant morphologic and textural character of each across-shore component, described in a landward to seaward sequence. Each component consists of a geomorphic form descriptor and a substrate descriptor:
FORM -> Bb, Csg, etc. TEXTURE

The primary geomorphic forms are initially described by a series of 12 codes:

A Anthropogenic E Coastal Bay, Lagoon, Estuary P Platform
B Beach I Inlet R River
C Cliff M Marsh S Bar/Trough
D Dune O Offshore Rocks T Delta

Each of these groups is then further modified by codes as indicated below.

Anthropogenic (A)
e causeway j jetty m marina t trench
f float g groin n boat ramp s seawall w wharf

Beach (B)
b berm i inclined slope s storm ridge
c washover channel n multiple intertidal bars t low-tide terrace
f beach face r single intertidal bar w washover

Cliff (C)
a active or erosional p passive c caves present

Dune (D)
b blowout f foredune s ridge and swale
d stabilized r random form l longitudinal

Coastal Bay, Lagoon, Estuary (E)
b enclosed bay c estuary

Inlet (I)
e ephemeral a opening fixed by offshore structures
s stable f flood-tidal delta l ebb-tidal delta

Marsh (M)
c tidal creek

Offshore Rocks (O)
e intertidal reef s sea stack
r rock outcrop (>2 m above M.S.L. and <10 m in width or length)

Platform (P)
b high-tide platform f horizontal t terraced
l low-tide platform r ramp i irregular

River (R)
b braided m multiple s single channel

Bar/Trough (S) subtidal r with rip channels

Delta (T)
c channel m multiple channels p delta plain
f fan s single channel v crevasse
l levee

Substrate type or sediment texture are described in detail by a series of codes. The sediments or materials of the shore are initially coded into one of four groups:
A anthropogenic materials C clastic sediments
B biogenic sediments R bedrock

Each of these groups is then further modified by texture or composition as indicated below.

Anthropogenic materials (A)
a metal n concrete (solid) w bark or wood debris
d debris, rubble t logs
e concrete (individually formed) u wood (structural; e.g., pilings or boards)

Biogenic sediments (B)
i trees or wood particles o organic litter
s shell hash (with a texture as described below)

Clastic sediments (C)
b boulder s sand m mud
c cobble s silt g gravel
p pebble f clay r rubble

Bedrock (R)
i igneous m metamorphic s sedimentary

Where more than one texture is present in an across-shore component, several substrates or textures may be indicated. Where one texture physically overlies another, it is indicated by a slash (e.g., Cs/Rs, and over rock). Where several mutually exclusive textures occur within a component, such as rock outcrops within a sand beach, a colon is used to indicate that association (e.g., Cs:R).

Prepared for
Minerals Management Service
Pacific Outer Continental Shelf Region
Los Angeles, California
by
Woodward-Clyde Consultants
Environmental Systems Division
San Francisco, California
November, 1982

M.M.S. Map# 77
Sheet 2 of 2